

WHAT TO DO WITH EQUINE FLU?

Steps to take when there is a case in your area or yard



REDUCE THE RISK

If your horse is fully vaccinated (completed a primary flu vaccine course and appropriately boosted), they should have good protection. If your horse encounters flu, clinical signs will typically be much less severe; your horse will get better faster and spread less virus, meaning other horses will be less likely to get sick.

Flu commonly occurs shortly after the arrival of new horses onto a premises*. Yards should have protocols for quarantining new arrivals for at least two weeks (ideally three weeks) before they mix with resident horses. Before arrival, confirm the new horse is vaccinated against flu (at least had its primary course, 3 injections, as a minimum) and discuss other infectious disease screening tests with your vet. Wash your hands between handling different horses and use designated equipment for each horse.

Equine events are good places for the circulation of infectious diseases. When away from the yard, take your own equipment, including bucket/s and water. Avoid communal areas and contact with other horses. Disinfect all equipment, including your trailer, when you return to your yard.

*[Whitlock et al., 2026](#) - *Equine influenza surveillance sheds light on sources and seasonal risk in the United Kingdom*

SPOTTING THE SIGNS

Flu is caused by a highly contagious virus and is transmitted by respiratory droplets through direct horse-to-horse contact and through the air with coughing.



One of the notable features of flu is the very quick spread of signs in a group of horses. The virus can spread large distances in the air, which is one reason why it is very important to control outbreaks as early as possible. Equipment and personnel (hands, clothing, etc) can also indirectly transmit the virus.

SIGNS OF FLU CAN INCLUDE:

- **Nasal discharge**
- **Coughing** - dry/harsh/hacking
- **Fever** (≥ 38.5 C)
- **Lethargy/poor performance**
- **Loss of appetite**



SEEK VETERINARY ADVICE AS SOON AS POSSIBLE IF YOU HAVE CONCERNS ABOUT YOUR HORSE



BE PREPARED TO ISOLATE

For flu, use a completely separate stable with no shared airspace, ideally at least 25 metres from other horses. Use separate equipment, handlers (or protective clothing, gloves, separate boots, and care for the isolated horse after all other horses), and muck heap when dealing with an isolated horse. Temporary isolation can be set up by moving other horses away from the stable area/block and using it just for the isolated horse.

Taping off the area and using clear signage makes others aware to avoid the area, with disinfectant foot dip and handwashing at the entrance/exit.



SUSPECT FLU?



CALL YOUR VET

They can take a swab sample from your horse's nose and throat to confirm if your horse has flu. If your horse has been sick for a while before you call the vet, it can be harder to diagnose your horse correctly. Swabs are best taken early on in the course of the infection.

Eligible samples can be tested for free through the equine influenza surveillance scheme. This scheme is kindly supported by the Horserace Betting Levy Board. If your vet hasn't signed up for the scheme, ask them to contact us.

Making a diagnosis helps to ensure that the best supportive care is given to your horse and that the correct preventative measures are put in place to stop the virus from spreading. Not only will you get a diagnosis, but you are also helping to monitor equine influenza activity in the UK. This allows us to study how the virus is evolving and helps us ensure the right strains are recommended for inclusion in future equine flu vaccines. This means that the most effective and up-to-date vaccines are produced to protect your horse from equine flu.

POST DIAGNOSIS STEPS

Your vet will advise you on treatment for the horse that has flu. As with human flu, rest, supportive care and anti-inflammatories are usually all that is needed and with time, your horse will feel better.

Vaccinated horses that have no signs can still carry and spread the virus, but the amount of virus shed should hopefully be at a lower level and for a shorter period of time. These clinically healthy horses should still follow the precautionary measures put in place. Measures to prevent the spread of flu are yard specific and tailor-made by your vet, with assistance from veterinary epidemiologists at EIDS.



RESEARCH UPDATE

Research by EIDS analysed surveillance data from laboratory-confirmed outbreaks of flu occurring in the UK from 2020 to 2024 which was characterised by a **notable October to December risk window and strong links to horse movements**, with spread of disease commonly amplified where vaccination and quarantine were suboptimal.

Read the full paper here: <https://doi.org/10.1002/evj.70156>

TAKE HOME MESSAGES



Infected horses will spread the virus through coughing and typically shed the virus for up to ten days, but this can be longer.



Infected horses should be isolated until your vet advises they are no longer infected.



All horse movements on and off the yard should be stopped.



Monitor all horses for clinical signs and record their rectal temperatures daily; they should be less than 38.5° C.



Ensure vaccines are up to date and discuss with your vet if it has been greater than six months since your horses last vaccine, as a booster may be indicated.



EIDS

Your vet can call EIDS through our free advice service for outbreak management advice that will be tailored to your horse and yard.

